

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 070 389**  
**A2**

(12)

# EUROPEAN PATENT APPLICATION

(21) Application number: 82105237.0

(22) Date of filing: 15.06.82

(51) Int. Cl.<sup>3</sup>: **C 07 D 471/04**, **C 07 D 513/04**,  
**A 01 N 43/50**, **C 07 D 211/60**,  
**C 07 D 279/12**  
//  
(C07D471/04, 235/00,  
221/00), (C07D513/04,  
279/00, 235/00)

(30) Priority: 16.08.81 JP 93315/81  
04.12.81 JP 196035/81  
04.12.81 JP 196036/81

(43) Date of publication of application: 26.01.83  
Bulletin 83/4

(84) Designated Contracting States: **CH DE FR GB IT LI NL**


(71) Applicant: **SUMITOMO CHEMICAL COMPANY, LIMITED**,  
15 Kitahama 5-chome Higashi-ku, Osaka-shi Osaka-fu  
(JP)

(72) Inventor: Nagano, Eiki, 4-1-401, Ryodo-cho Nishinomiya,  
Hyogo (JP)  
Inventor: Hashimoto, Shunichi, 2-10-4-442,  
Sonehigashi-machi, Toyonaka Osaka (JP)  
Inventor: Yoshida, Ryo, 5-19, Azajizoyama Higashiuneno,  
Kawanishi Osaka (JP)  
Inventor: Matsumoto, Hiroshi, 2-10-2-221,  
Sonehigashi-machi, Toyonaka Osaka (JP)  
Inventor: Kamoshita, Katsuzo, 2-3-11, Kofudai  
Toyono-cho, Toyono-gun Osaka (JP)

(74) Representative: **Vossius Vossius Tauchner Heunemann**  
Rauh, Siebertstrasse 4 P.O. Box 86 07 67,  
D-8000 München 86 (DE)

(54) **2-(2-Fluoro-4-halo-5-substituted phenyl) hydantoins, and their production and use.**

(57) A herbicidal composition which comprises as an active ingredient a compound of the formula:



wherein X is a chlorine atom or a bromine atom, Y is  $-\text{CH}_2-$ ,  $-\text{S}-$  or  $-\text{SO}_2-$  and R is a  $\text{C}_1-\text{C}_4$  alkyl group, an allyl group or a propargyl group, and an inert carrier.

**EP 0 070 389 A2**


ACTORUM AG

**BEST AVAILABLE COPY**

2-(2-FLUORO-4-HALO-5-SUBSTITUTED PHENYL)HYDANTOINS,  
AND THEIR PRODUCTION AND USE

The present invention relates to 2-(2-fluoro-4-halo-5-substituted phenyl)hydantoins (hereinafter referred to as "hydantoin(s)"), and their production and use.

The said hydantoins are representable by the formula:



(I)

wherein X is a chlorine atom or a bromine atom, Y is  $-CH_2-$ ,  $-S-$  or  $-SO_2-$  and R is a  $C_1-C_4$  alkyl group, an allyl group or a propargyl group.

It is known that certain kinds of N-phenyl-hydantoins are effective as herbicides. For instance, the herbicidal use of 2-(4-chlorophenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine-1,3(2H,8aH)-dione, 2-(4-chloro-2-fluorophenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine-1,3(2H,8aH)-dione, 1,2-(4-chloro-2-fluorophenyl)-5,6,7,8-tetrahydro-1H-imidazo[5,1-c][1,4]thiadine-1,3(2H)-dione, 2-(4-chloro-2-fluorophenyl)-5,6,7,8-tetrahydro-1H-imidazo[5,1-c][1,4]thiadine-1,3(2H)-dione-7,7-dioxide, etc. is disclosed in British patent 1,503,244, U.S. patent 3,958,976, 4,179,276, etc. However, their herbicidal effect is still not necessarily satisfactory.


It has now been found that the hydantoins (I) show a strong herbicidal activity against a wide variety of weeds including Gramineae weeds, Cyperaceae weeds and broad-leaved weeds at small doses and do not produce any material phytotoxicity on various agricultural crops (i.e. corn, soybean, cotton, rice plant, wheat). Examples of Gramineae weeds against which the hydantoins (I) show a herbicidal activity are barnyardgrass (Echinochloa crus-galli), green foxtail (Setaria viridis), large crabgrass (Digitaria sanguinalis), Johnsongrass (Sorghum halepense), wild oat (Avena fatua), water foxtail (Alopecurus geniculatus), goosegrass (Eleusine indica), annual bluegrass (Poa annua), bermudagrass (Cynodon dactylon), quackgrass (Agropyron repens), etc. Examples of broad-leaved weeds are tall morningglory (Ipomoea purpurea), velvetleaf (Abutilon theophrasti), sicklepod (Cassia obtusifolia), wild sunflower (Helianthus annus), cocklebur (Xanthium pennsylvanicum), wild mustard (Brassica kaber), common purslane (Portulaca oleracea), jimsonweed (Datura stramonium), hemp sesbania (Sesbania exaltata), sun spurge (Euphorbia helioscopia), black nightshade (Solanum nigrum), prickly sida (Sida spinosa), common ragweed (Ambrosia artemisifolia), smartweed sp. (Polygonum sp.), redroot pigweed (Amaranthus retroflexus), bedstraw (Galium aparine), pineappleweed (Matricaria spp.), birdseye speedwell (Veronica persica), wild buckwheat (Polygonum convolvulus), beggarticks (Bidens spp.), common lambsquarters (Chenopodium album), bindweed (Calystegia japonica), monochoria

(Monochoria vaginalis), Dopatrium junceum, waterwort  
(Elatine triandra), false pimpernel (Lindernia procumbens),  
toothcup (Rotala indica), arrowhead (Sagittaria pygmaea),  
etc. Examples of Cyperaceae weeds are nutsedge sp. (Cyperus  
sp.), Cyperus rotundus, Cyperus esculentus, hardstem bulrush  
(Scirpus juncoides), nutsedge (Cyperus serotinus), water  
chestnut (Eleocharis kuroguwai), slender spikerush  
(Eleocharis acicularis), etc. Accordingly, the hydantoins  
(I) can be used as herbicides applicable to paddy field as  
well as agricultural plowed field. They are also useful as  
herbicides to be employed for orchard, tea garden, mulberry  
field, rubber plantation, forest, lawn, pasture, non-  
agricultural field, etc.

The hydantoins (I) can be produced by various  
procedures, among which typical examples are shown below:


Procedure A

The hydantoin (I) is obtainable by reacting a  
phenyl isocyanate of the formula:




(II)

wherein X and R are each as defined above with an azine of  
the formula:



(III)


wherein R' is an alkali metal atom or an alkyl group, and Y is as defined above in an inert solvent (e.g. water, toluene, hexane, benzene) at a temperature from room temperature (ca. 20°C) to the boiling temperature of the solvent to give an urea of the formula:



wherein X, Y, R and R' are each as defined above, followed by treatment of the latter with a mineral acid such as hydrochloric acid or an alkali metal alkoxide such as sodium methoxide for ring closure.

#### Procedure B

The hydantoin (I) is obtainable by reacting a hydroxyphenylhydantoin of the formula:



wherein X and Y are each as defined above with a halide of the formula:



wherein Z is a chlorine atom, a bromine atom or an iodine atom and R is as defined above in the presence of a base (e.g. potassium carbonate, sodium hydroxide, sodium hydride, sodium alkoxide) in an inert solvent (e.g. dimethylformamide, dimethylsulfoxide, acetonitrile, methylethylketone,


ethylene glycol monomethyl ether) at a temperature of 0 to 200°C. The molar ratio of the halide (V) and the base is preferred to be from 1.0 to 2.0.


Procedure C

The hydantoin (I:  $Y = -SO_2-$ ) is obtainable by oxidizing the corresponding hydantoin (I:  $Y = -S-$ ) with a peracid (e.g. peracetic acid, perbenzoic acid, m-chloro perbenzoic acid) in the presence of an inert solvent (e.g. methylene chloride, chloroform) at a temperature of -30 to 30°C.

The thus produced hydantoin (I) may be, when desired, purified by a per se conventional procedure such as recrystallization or column chromatography.

The production of the starting materials are summarized in the following scheme:






wherein X, Y and R are each as defined above.

Namely, the hydroxyphenyl isocyanate (X) may be produced from the phenol (VII) by nitrating the same, subjecting the resultant nitrophenol (VIII) to reduction and reacting the resulting aminophenol (IX) with phosgene. The thus produced hydroxyphenyl isocyanate (X) is then reacted with the azine (III) to give the hydroxyphenylhydantoin (V). Also, the phenyl isocyanate (II) may be produced from the nitrophenol (VIII) by subjecting the latter to alkylation, alkenylation or alkynylation, reducing the resultant nitrobenzene (XI) and reacting the resulting aminobenzene (XII) with phosgene.

Explaining the above conversions more in detail, the nitration of the phenol (VII) into the nitrophenol (VIII) may be accomplished by application of a per se conventional nitration procedure to the former. Usually, however, the indirect nitration which consists of the following three steps is favorable in achievement of the selective nitration at the desired position:





wherein X is as defined above. Thus, the phenol (VII) (Finger et al.: J.Am.Chem.Soc., 81, 94 (1959)) is converted into its alkali metal salt by treatment with an aqueous solution of an alkali metal hydroxide (e.g. sodium hydroxide, potassium hydroxide), and the resulting salt is reacted with an alkyl haloformate such as methyl chloroformate in water at a temperature of 0 to 10°C. The thus prepared carbonic ester (XIII) is nitrated with a mixture of conc. sulfuric acid and conc. nitric acid at room temperature. Then, the nitrobenzene (XIV) thus obtained is hydrolyzed with an aqueous alkaline solution such as an aqueous sodium hydroxide solution at a temperature of 40 to 80°C to give the nitrophenol (VIII).

Reduction of the nitrophenol (VIII) into the aminophenol (IX) may be carried out by a per se conventional procedure for reduction of a nitro group on an aromatic ring to an amino group. Examples of such procedure is reduction with sodium sulfide or iron powder, catalytic reduction, etc. Preferred are reduction with gaseous hydrogen in the presence of platinum dioxide as a catalyst in an amount of 1/10 to 1/100 mol to 1 mol of the nitrophenol (VIII) at room temperature under atmospheric pressure, reduction of the nitrophenol (VIII) with iron powder (e.g. reductive iron, electrolytic iron) in an amount of 2.0 to 5.0 mol to 1 mol of the nitrophenol (VIII) in acetic acid at a temperature of 80 to 120°C, etc.

The reaction of the aminophenol (IX) with phosgene

in an inert solvent (e.g. anhydrous ethyl acetate) gives the hydroxyphenyl isocyanate (X).

The hydroxyphenyl isocyanate (X) is reacted with the azine (III) at room temperature in an inert solvent (e.g. benzene, toluene, xylene), if necessary, by treatment with a mineral acid to give the hydroxyphenylhydantoin (V).

Alternatively, the alkylation, alkenylation or alkynylation for conversion of the nitrophenol (VIII) to nitrobenzene (XI) may be carried out by treatment of the former with an alkali metal carbonate (e.g. potassium carbonate), an alkali metal hydride (e.g. sodium hydride) or an alkali metal alkoxide (e.g. sodium methoxide) and reacting the resultant alkali metal salt with the halide (VI) in a polar solvent (e.g. water, dimethylformamide, acetonitrile, acetone, dimethylsulfoxide), usually at a temperature of 10 to 200°C, preferably of 30 to 100°C. The use of a phase transfer catalyst such as tetrabutylammonium bromide is favorable for smooth accomplishment of the reaction.

Reduction of the nitrobenzene (XI) to the aminobenzene (XII) may be achieved in various procedures. When, for instance, R in the compound (XI) is C<sub>1</sub>-C<sub>4</sub> alkyl, there may be adopted a per se conventional reduction procedure for converting a nitro group into an amino group wherein a reducing agent such as sodium sulfide or iron powder or catalytic reduction is employed. One of typical procedures comprises introduction of a 3 molar amount of

hydrogen into a reaction system comprising one molar amount of the compound (XI) and a 1/10 to 1/100 molar amount of platinum dioxide at room temperature under atmospheric pressure. Another typical procedure comprises admixing an acetic acid solution containing one molar amount of the compound (XI) with a 5 % acetic acid solution containing a 2.5 to 5.0 molar amount of iron powder such as reductive iron or electrolytic iron and effecting the reaction at a temperature of 80 to 100°C. When R in the compound (XI) is propargyl or allyl, there may be adopted reduction with iron powder. For instance, an acetic acid solution containing one molar amount of the compound (XI) may be admixed with a 5 % acetic acid solution containing 2.5 to 5.0 molar amount of iron powder such as reductive iron or electrolytic iron at a temperature of 80 to 120°C, preferably of 90 to 110°C, for a period of 0.5 to 5 hours.

The aminobenzene (XII) is converted into the phenyl isocyanate (II) by reacting the former with phosgene in an inert solvent (e.g. toluene, benzene, ethyl acetate, tetrahydrofuran, 1,4-dioxane).

Practical and presently preferred embodiments of the production of the objective hydantoins (I) as well as the intermediary compounds are illustratively shown below:

Example 1

A solution of 4-chloro-2-fluoro-5-isopropoxyphenyl isocyanate (2.3 g) in chlorobenzene (10 ml) was added to a 2 % aqueous solution of sodium hydroxide (25 ml) containing

pipecolic acid (1.3 g), and the mixture was stirred at room temperature (ca. 20°C) for 3 hours. After allowed to stand overnight, the precipitated crystals were collected by filtration and washed with ether. The crystals were combined with water (20 ml) and adjusted to pH 2 or less with hydrochloric acid, followed by reflux for 3 hours. After allowed to cool, the mixture was extracted with ether, dried and concentrated. The residue was purified by silica gel chromatography to give 0.6 g of 2-(4-chloro-2-fluoro-5-isopropoxyphenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine-1,3(2H,8aH)-dione (Compound No. 7). M.P. 60.5 - 64.5°C.

NMR (CDCl<sub>3</sub>),  $\delta$  ppm: 1.4 (6H, d, J=6Hz), 4.2 - 4.6 (1H, m), 6.8 (1H, d, J=6Hz), 7.2 (1H, d, J=10Hz).

#### Example 2

To a solution of 2-(4-chloro-2-fluoro-5-hydroxyphenyl)-5,6,7,8-tetrahydroimidazo[1,5a]pyridine-1,3(2H,8aH)-dione (1 g) in dimethylformamide (10 ml), there were added anhydrous potassium carbonate (0.3 g) and propargyl bromide (1 g), and the mixture was stirred at 60 - 70°C for 3 hours. After allowed to cool, water was added thereto, followed by extraction with ether. The ether layer was washed with water, dried and concentrated. The residue was recrystallized from ether and petroleum ether to give 0.5 g of 2-[4-chloro-2-fluoro-5-(1-propynyloxy)phenyl]-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine-1,3(2H,8aH)-dione (Compound No. 10). M.P. 124 - 125°C.

NMR (CDCl<sub>3</sub>)  $\delta$  ppm: 2.6 (1H, t, J=3Hz), 3.0 (1H,

t like m), 4.72 (2H, d, J=3Hz), 7.0 (1H, d, J=6Hz), 7.25 (1H, d, J=10Hz).

### Example 3

To a solution of 2-methoxycarbonyl-4-thiomorpholine (6 g) in n-hexane (60 ml), there was added a catalytic amount of triethylamine, and 4-chloro-2-fluoro-5-methoxyphenyl isocyanate (6.9 g) was portionwise added thereto at room temperature, whereby an insoluble oily substance was solidified. After removal of n-hexane by decantation, methanol (60 ml) and a catalytic amount of sodium methoxide were added thereto, and the mixture was refluxed for 1 hour. The reaction mixture was concentrated, washed with water and extracted with ether. The ether extract was washed with water, dried over anhydrous magnesium sulfate and concentrated. The residue was purified by silica gel chromatography to give 2.6 g of 1,2-(4-chloro-2-fluoro-5-methoxyphenyl)-5,6,8,8a-tetrahydro-1H-imidazo[5,1-c][1,4]thiazine-1,3(2H)-dione (Compound No. 13). M.P. 121 - 123°C.

IR  $\nu_{\max}$  ( $\text{cm}^{-1}$ ): 1780, 1720.

### Example 4

1,2-(4-Chloro-2-fluoro-5-methoxyphenyl)-5,6,8,8a-tetrahydro-1H-imidazo[5,1-c][1,4]thiazine-1,3(2H)-dione (1.6 g) as produced in Example 3 was dissolved in methylene chloride (30 ml), and a 70 % m-chloroperbenzoic acid (2.2 g) was added thereto. The reaction was carried out at a temperature of -10 - 0°C for 6 hours. Upon confirming


no presence of peracid by an iodo-starch paper, the reaction mixture was washed with sodium bicarbonate solution. The organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was recrystallized from ether and petroleum ether to give 0.9 g of 1,2-(2-fluoro-4-chloro-5-methoxyphenyl)-5,6,8,8a-tetrahydro-1H-imidazo[5,1-c][1,4]-thiazine-1,3(2H)-dione-7,7-oxide (Compound No. 14). M.P. 195 - 196°C.

Example 5

1,2-(2-Fluoro-4-chloro-5-hydroxyphenyl)-5,6,8,8a-tetrahydro-1H-imidazo[5,1-c][1,4]thiazine-1,3(2H)-dione-7,7-oxide (1.3 g), potassium carbonate (0.28 g) and propargyl bromide (0.5 g) were dissolved in dimethylformamide (20 ml), and the mixture was stirred at 50 - 61°C for 2.5 hours. After allowed to cool, the resultant mixture was extracted with ethyl acetate. The extract was washed with water, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography to give 1.0 g of 1,2-(2-fluoro-4-chloro-5-propargyloxyphenyl)-5,6,8,8a-tetrahydro-1H-imidazo[5,1-c][1,4]thiazine-1,3(2H)-dione-7,7-oxide (Compound No. 25). M.P. 178 - 181°C.

Examples of the hydantoin (I) produced in the same manner as above are shown in Table 1.

Table 1



Compound No.	X	Y	R	Physical property
1	Cl	CH <sub>2</sub>	-CH <sub>3</sub>	M.P. 162 - 163°C
2	Br	CH <sub>2</sub>	-CH <sub>3</sub>	M.P. 170.2 - 171.8°C
3	Cl	CH <sub>2</sub>	-CH <sub>2</sub> CH <sub>3</sub>	Glassy
4	Br	CH <sub>2</sub>	-CH <sub>2</sub> CH <sub>3</sub>	M.P. 131.5 - 133°C
5	Cl	CH <sub>2</sub>	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	Glassy
6	Br	CH <sub>2</sub>	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	Glassy
7	Cl	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>	M.P. 60.5 - 64.5°C
8	Br	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>	Glassy
9	Cl	CH <sub>2</sub>	-CH <sub>2</sub> CH=CH <sub>2</sub>	M.P. 111 - 112°C
10	Cl	CH <sub>2</sub>	-CH <sub>2</sub> C≡CH	M.P. 124 - 125°C
11	Br	CH <sub>2</sub>	-CH <sub>2</sub> C≡CH	M.P. 149 - 150°C
12	Cl	CH <sub>2</sub>	-(CH <sub>2</sub> ) <sub>3</sub> CH <sub>3</sub>	Glassy
13	Cl	S	-CH <sub>3</sub>	M.P. 121 - 123°C
14	Cl	SO <sub>2</sub>	-CH <sub>3</sub>	M.P. 195 - 196°C
15	Cl	S	-CH <sub>2</sub> CH <sub>3</sub>	M.P. 125 - 126.5°C

(Continued)

Compound No.	X	Y	R	Physical property
16	Cl	SO <sub>2</sub>	-CH <sub>2</sub> CH <sub>3</sub>	M.P. 218 - 220°C
17	Br	S	-CH <sub>2</sub> CH <sub>3</sub>	n <sub>D</sub> <sup>24.5</sup> 1.5623
18	Br	S	-CH <sub>2</sub> CH <sub>3</sub>	M.P. 93 - 94.5°C
19	Cl	S	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	n <sub>D</sub> <sup>24.0</sup> 1.5540
20	Cl	SO <sub>2</sub>	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	M.P. 107 - 110°C
21	Cl	S	-CH(CH <sub>3</sub> ) <sub>2</sub>	Glassy
22	Cl	SO <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>	M.P. 69.4°C
23	Cl	S	-CH(CH <sub>3</sub> )CH <sub>2</sub> CH <sub>3</sub>	M.P. 143 - 144°C
24	Cl	SO <sub>2</sub>	-CH(CH <sub>3</sub> )CH <sub>2</sub> CH <sub>3</sub>	M.P. 154 - 155°C
25	Cl	SO <sub>2</sub>	-CH <sub>2</sub> C≡CH	M.P. 178 - 181°C


Example 6

A solution of 4-chloro-2-fluoro-5-isopropoxy-aniline (30 g) in toluene (100 ml) was added to a 1 M phosgene/toluene solution (500 ml) at room temperature (ca. 20°C), followed by heating under reflux. The mixture was concentrated under reduced pressure, and the residue was distilled to give 26 g of 4-chloro-2-fluoro-5-isopropoxy-phenyl isocyanate as pale yellow crystals. M.P., 36 - 37°C. B.P., 90 - 91°C/3 mmHg.

IR  $\nu_{\max}$  (cm<sup>-1</sup>): 2240.

Some examples of the phenyl isocyanate (II) produced in the same manner as above are shown in Table 2.



Table 2

X	R	Physical property
Cl	CH <sub>3</sub> -	M.P. 42 - 44.5°C
Cl	n-C <sub>3</sub> H <sub>7</sub> -	M.P. 43 - 44°C
Cl	iso-C <sub>3</sub> H <sub>7</sub> -	M.P. 36 - 37°C
Br	C <sub>2</sub> H <sub>5</sub> -	M.P. 35 - 36.5°C
Cl	CH <sub>2</sub> =CHCH <sub>2</sub> -	B.P. 107°C/3 mmHg; n <sub>D</sub> <sup>16</sup> 1.5481
Cl	CH≡CCH <sub>2</sub> -	M.P. 61.5 - 62.5°C
Cl	sec-C <sub>4</sub> H <sub>9</sub> -	B.P. 102°C/1 mmHg; n <sub>D</sub> <sup>22.0</sup> 1.5200

Example 7

A mixture of 4-chloro-2-fluoro-5-isopropoxy-nitrobenzene (13.5 g) and platinum dioxide (0.4 g) in ethanol (300 ml) was subjected to catalytic reduction with hydrogen at room temperature (ca. 20°C) under atmospheric pressure, whereby a designed amount of hydrogen was absorbed. The resultant mixture was filtered to remove insoluble materials, and the filtrate was concentrated. The residue was subjected to purification by silica gel chromatography to give 5.6 g of 4-chloro-2-fluoro-5-isopropoxyphenylaniline. n<sub>D</sub><sup>24.5</sup> 1.5360.

NMR (CDCl<sub>3</sub>) δ (ppm): 1.3 (6H, d, J=6Hz), 3.7 (2H, m, J=1.5Hz), 4.35 (1H, q, J=6Hz), 6.45 (1H, d, J=7Hz), 7.1 (1H, d, J=10Hz).

IR  $\nu_{\max}$  ( $\text{cm}^{-1}$ ): 3450, 3550.

Example 8


A suspension of electrolytic iron powder (3.5 g) in a 5 % aqueous acetic acid solution (5 ml) was heated to 90°C, and a solution of 4-chloro-2-fluoro-5-(2-propynyloxy)-nitrobenzene (5.7 g) in acetic acid (40 ml) was dropwise added thereto at the same temperature. The resultant mixture was stirred at 90 - 105°C for 1 hour and allowed to cool to room temperature. Water (200 ml) was added thereto. Insoluble materials were filtered off, and the filtrate was neutralized, followed by extraction with ethyl acetate. The extract was dried over anhydrous magnesium sulfate and concentrated. The residue was washed with petroleum ether and carbon tetrachloride to give 3.6 g of 4-chloro-2-fluoro-5-(2-propynyloxy)aniline. M.P. 61.0 - 61.5°C.

NMR ( $\text{CDCl}_3$ )  $\delta$  (ppm): 2.5 (1H, t,  $J=2\text{Hz}$ ), 3.4 - 4.2 (2H, m,  $J=16\text{Hz}$ ), 4.15 (2H, d,  $J=2\text{Hz}$ ), 6.5 (1H, d,  $J=8\text{Hz}$ ), 6.95 (1H, d,  $J=10\text{Hz}$ ).

IR  $\nu_{\max}$  ( $\text{cm}^{-1}$ ): 3460, 3360, 3280, 2100.

Some examples of the aminobenzene (XII) produced in the same manner as above are shown in Table 3.

---

Table 3

X	R	Physical property
Cl	C <sub>2</sub> H <sub>5</sub> -	n <sub>D</sub> <sup>24.5</sup> 1.5503
Br	C <sub>2</sub> H <sub>5</sub> -	n <sub>D</sub> <sup>25.0</sup> 1.5680
Cl	n-C <sub>3</sub> H <sub>7</sub> -	n <sub>D</sub> <sup>24.5</sup> 1.5386
Br	n-C <sub>3</sub> H <sub>7</sub> -	n <sub>D</sub> <sup>26.0</sup> 1.5618
Cl	iso-C <sub>3</sub> H <sub>7</sub> -	n <sub>D</sub> <sup>24.5</sup> 1.5360
Br	iso-C <sub>3</sub> H <sub>7</sub> -	n <sub>D</sub> <sup>25.0</sup> 1.5547
Cl	CH <sub>2</sub> =CHCH <sub>2</sub> -	n <sub>D</sub> <sup>19</sup> 1.5598
Cl	CH≡CCH <sub>2</sub> -	M.P. 61.0 - 61.5°C
Cl	CH≡C-CH-   CH <sub>3</sub>	M.P. 67.0 - 68°C

Example 9

To a solution of 2-chloro-4-fluoro-5-nitrophenol (19.1 g) in acetonitrile (100 ml), there was added anhydrous potassium carbonate (8.g). After stirring at room temperature (ca. 20°C) for several minutes, isopropyl iodide (25 g) was added thereto, and the resultant mixture was heated under reflux for 3 hours. After allowed to cool to room temperature (ca. 20°C), water was added thereto, and the reaction mixture was extracted with ether. The ether extract was washed with a 5 % aqueous sodium hydroxide solution and water in order, dried over anhydrous magnesium

sulfate and filtered. The filtrate was concentrated under reduced pressure, and the residue was recrystallized from ethanol to give 13.5 g of 4-chloro-2-fluoro-5-isopropoxy-nitrobenzene. M.P., 61.3 - 62.4°C.

NMR (CDCl<sub>3</sub>)  $\delta$  (ppm): 1.42 (6H, d, J=7Hz), 4.3 - 4.8 (1H, m), 7.28 (1H, d, J=10Hz), 7.48 (1H, d, J=6Hz).

Some examples of the nitrobenzene (XI) produced in the same manner as above are shown in Table 4.





Table 4



X	R	Physical property
Cl	-CH <sub>3</sub>	M.P. 67.5 - 69.8°C
Br	-CH <sub>3</sub>	M.P. 72.2°C
Cl	-CH <sub>2</sub> CH <sub>3</sub>	M.P. 47 - 48°C
Br	-CH <sub>2</sub> CH <sub>3</sub>	M.P. 46 - 46.5°C
Cl	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	M.P. 46 - 47°C
Br	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	M.P. 46.8 - 47.4°C
Cl	-CH(CH <sub>3</sub> ) <sub>2</sub>	M.P. 61.3 - 62.4°C
Br	-CH(CH <sub>3</sub> ) <sub>2</sub>	M.P. 65.5 - 66.5°C
Cl	-CH(CH <sub>3</sub> )CH <sub>2</sub> CH <sub>3</sub>	M.P. 59.6 - 60.6°C
Cl	-CH <sub>2</sub> CH=CH <sub>2</sub>	n <sub>D</sub> <sup>17.0</sup> 1.5601
Cl	-CHCH=CH <sub>2</sub>   CH <sub>3</sub>	M.P. 41.0 - 41.5°C
Cl	-CH <sub>2</sub> C≡CH	M.P. 88 - 89°C
Cl	-CHC≡CH   CH <sub>3</sub>	M.P. 87 - 88°C

Example 10

2-Chloro-4-fluorophenol (83.4 g) was added to a solution of sodium hydroxide (27.7 g) in water (450 ml), and methyl chloroformate (69.2 g) was dropwise added thereto at a temperature below 10°C. Precipitated crystals were collected by filtration and washed with water to give methyl

(2-chloro-4-fluorophenyl)formate (134.8 g). M.P., 69 - 71°C.

Methyl (2-chloro-4-fluorophenyl)formate (134.8 g) obtained above was suspended in conc. sulfuric acid (50 ml). To the suspension, a mixture of conc. sulfuric acid (50 ml) and conc. nitric acid (50 ml) was added at about 30°C, and the mixture was stirred at this temperature for 1 hour. The reaction mixture was poured into ice water, and precipitated crystals were collected and washed with water to give methyl (2-chloro-4-fluoro-5-nitrophenyl)formate (143 g). M.P., 50 - 55°C.

The product obtained as above was combined with sodium hydroxide (27 g) and water (300 ml), and the resultant mixture was refluxed for 4 hours. Precipitated insoluble materials were filtered using a celite, and the filtrate was acidified with conc. hydrochloric acid. Precipitated crystals were collected by filtration and washed with water to obtain 76.3 g of 2-chloro-4-fluoro-5-nitrophenol. M.P., 106 - 107°C.

NMR ( $\text{CDCl}_3$ ,  $\text{D}_6$ -DMSO)  $\delta$  (ppm): 7.25 (1H, d,  $J=10\text{Hz}$ ), 7.64 (1H, d,  $J=6\text{Hz}$ ).

IR  $\nu_{\text{max}}^{\text{nujol}}$  ( $\text{cm}^{-1}$ ): 3370.

#### Example 11

To a solution of 2-bromo-4-fluorophenol (28 g) in a solution of sodium hydroxide (7 g) in water (100 ml), methyl chloroformate was dropwise added thereto at a temperature below 10°C. The produced crystals were

collected by filtration and washed with water to give methyl 2-bromo-4-fluorophenylformate (41 g). M.P., 80.7°C.

The above product was suspended in conc. sulfuric acid (13 ml), a mixture of conc. sulfuric acid (13 ml) and conc. nitric acid (13 ml) was dropwise added thereto at about 30°C, and the resultant mixture of stirred for 30 minutes. The reaction mixture was poured into ice water. The produced crystals were collected by filtration and washed with water to give methyl 2-bromo-4-fluoro-5-nitroformate (38.3 g) as yellow crystals. M.P., 63.5 - 64.5°C.

The thus obtained product was admixed with sodium hydroxide (6.2 g) and water (100 ml) and heated under reflux for 3 hours. The insoluble materials were eliminated by filtration, and the filtrate was acidified with hydrochloric acid. The precipitated crystals were collected by filtration and washed with water to give 25 g of 2-bromo-4-fluoro-5-nitrophenol. M.P., 126 - 127°C.

NMR ( $\text{CDCl}_3$ ,  $\text{D}_6$ -DMSO)  $\delta$  (ppm): 7.42 (1H, d,  $J=10\text{Hz}$ ), 7.65 (1H, d,  $J=6\text{Hz}$ ).

IR  $\nu_{\text{max}}^{\text{nujol}}$  ( $\text{cm}^{-1}$ ): 3450.

#### Example 12


4-Bromo-2-fluoro-5-hydroxyphenyl isocyanate (3 g) was dissolved in chlorobenzene (5 ml), and a solution of pipecolic acid (1.7 g) and sodium hydroxide (0.56 g) in water (5 ml) was added thereto, followed by stirring at room temperature (ca. 20°C) overnight. The aqueous layer was washed with ether, made acidic with an aqueous hydrochloric

acid solution and heated under reflux for 2 hours. After allowed to cool, the resulting mixture was extracted with methylene chloride (300 ml). The extract was washed with sodium bicarbonate solution, dried and concentrated. The residue was recrystallized from ether to give 1.75 g of 2-(4-bromo-2-fluoro-5-hydroxyphenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine-1,3(2H,8aH)-dione. M.P., 196 - 199°C.

IR  $\nu_{\text{max}}$  ( $\text{cm}^{-1}$ ): 3400, 1760, 1700.

Some examples of the hydroxyhydantoin (V) produced in the same manner as above are shown in Table 5.

Table 5



X	Y	Physical property
Cl	-CH <sub>2</sub> -	M.P. 193.5 - 194°C
Br	-CH <sub>2</sub> -	M.P. 196 - 199°C
Cl	-S-	M.P. 207.5 - 209.5°C
Cl	-SO <sub>2</sub> -	Glassy

Example 13

Ethyl pipecolate hydrochloride (4.64 g) was treated with a 1 % sodium hydroxide solution (100 ml), followed by extraction with toluene. The toluene layer was washed with a saturated sodium chloride solution, 4-chloro-2-fluoro-(1-methylethoxy)phenyl isocyanate (5 g) was drop-




wise added thereto, and the resultant mixture was stirred for 4 hours. Toluene was removed under reduced pressure to give 5.93 g of ethyl N-[4-chloro-2-fluoro-5-(1-methylethoxy)phenylcarbamoyl]pipercolate as a glassy substance.

NMR ( $\text{CDCl}_3$ )  $\delta$  ppm: 1.25 (3H, t), 1.32 (6H, d), 4.2 (2H, q), 7.82 (1H, d).

Some examples of the urea (IV) produced in the same manner as above are shown in Table 6.

Table 6



X	Y	R	R'	Physical property
Cl	$\text{CH}_2$	iso- $\text{C}_3\text{H}_7$	Na	M.P. 59.5 - 65°C
Cl	$\text{CH}_2$	iao- $\text{C}_3\text{H}_7$	$\text{C}_2\text{H}_5$	Glassy
Cl	S	iso- $\text{C}_3\text{H}_7$	$\text{C}_2\text{H}_5$	M.P. 114 - 116°C

In the practical usage of the hydantoins (I), they may be applied as such or in any preparation form such as dusts, granules, wettable powders, emulsifiable concentrates or suspensions.

The concentration of the active ingredient in such preparation form is usually within a range of 0.1 to 80 % by weight. However, the upper and lower range may be also effective depending upon purposes.

For formulation of those preparations, a solid or liquid carrier or diluent may be used. As the solid carrier

or diluent, there may be employed mineral powders (e.g. kaolin, bentonite, montmorillonite, talc, diatomaceous earth, mica, vermiculite, gypsum, calcium carbonate, apatite, synthetic hydrous silicate), vegetable powders (e.g. soybean powder, wheat flour, wooden powder, tobacco powder, starch, crystalline cellulose), high molecular weight compounds (e.g. petroleum resin, polyvinyl chloride, dammar gum, ketone resin), alumina, wax, etc.

As the liquid carrier or diluent, there may be employed alcohols (e.g. methanol, ethanol, ethylene glycol, benzyl alcohol), aromatic hydrocarbons (e.g. toluene, benzene, xylene, methylnaphthalene), halogenated hydrocarbons (e.g. chloroform, carbon tetrachloride, monochlorobenzene), ethers (e.g. dioxane, tetrahydrofuran), ketones (e.g. acetone, methylethylketone, cyclohexanone), esters (e.g. ethyl acetate, butyl acetate, ethylene glycol acetate), acid amides (e.g. dimethylformamide), nitriles (e.g. acetonitrile), ether alcohols (e.g. ethylene glycol ethyl ether), water, etc.

A surface active agent used for emulsification, dispersion or spreading may be any of the non-ionic, anionic, cationic and amphoteric type of agents. Examples of the surface active agent include polyoxyethylene alkyl ethers, polyoxyethylene alkylaryl ethers, polyoxyethylene fatty acid esters, sorbitan fatty acid esters, polyoxyethylene sorbitan fatty acid esters, oxyethyleneoxypropylene polymers, polyoxyethylene alkyl phosphates, fatty acid

salts, alkyl sulfates, alkyl sulfonates, alkylaryl sulfonates, alkyl phosphates, polyoxyethylene alkyl sulfates, quaternary ammonium salts and the like. If necessary, gelatin, casein, sodium alginate, starch, agar, polyvinyl alcohol, ligninsulfonates, isopropyl acid phosphate or the like may be used as an auxiliary agent.

The hydantoins (I) of the invention may be used together with other herbicides to improve their activity as herbicides, and in some cases, a synergistic effect can be expected. As the other herbicides, there may be exemplified phenoxy series herbicides such as 2,4-dichlorophenoxyacetic acid, 2-methyl-4-chlorophenoxybutyric acid and 2-methyl-4-chlorophenoxyacetic acid (including esters and salts thereof); diphenyl ether series herbicides such as 2,4-dichlorophenyl-4'-nitrophenyl ether, 2,4,6-trichlorophenyl-4'-nitrophenyl ether, 2-chloro-4-trifluoromethylphenyl-3'-ethoxy-4'-nitrophenyl ether, 2,4-dichlorophenyl-4'-nitro-3'-methoxyphenyl ether and 2,4-dichlorophenyl-3'-methoxycarbonyl-4'-nitrophenyl ether; triazine series herbicides such as 2-chloro-4,6-bisethylamino-1,3,5-triazine, 2-chloro-4-ethylamino-6-isopropylamino-1,3,5-triazine, 2-methylthio-4,6-bisethylamino-1,3,5-triazine, 2-methylthio-4,6-bis-isopropylamino-1,3,5-triazine and 4-amino-6-t-butyl-3-methylthio-1,2,4-triazin-5-one; urea series herbicides such as 3-(3,4-dichlorophenyl)-1,1-dimethylurea, 3-(3,4-dichlorophenyl)-1-methoxy-1-methylurea, 1-(2,2-dimethylbenzyl)-3-p-tolylurea and 1,1-dimethyl-3-(3-trifluoromethylphenyl)urea;

carbamate series herbicides such as isopropyl N-(3-chlorophenyl)carbamate and methyl N-(3,4-dichlorophenyl)carbamate; thiolcarbamate series herbicides such as S-(4-chlorobenzyl)-N,N-diethylthiolcarbamate, S-ethyl-N,N-hexamethylenethiolcarbamate, S-ethyl-N,N-diisobutylthiolcarbamate, S-ethyl-N,N-di-n-propylthiolcarbamate and S-n-propyl-N,N-di-n-propylthiolcarbamate; anilide series herbicides such as 3,4-dichloropropionanilide, N-methoxymethyl-2,6-diethyl- $\alpha$ -chloroacetanilide, 2-chloro-2',6'-diethyl-N-(butoxymethyl)-acetanilide, 2-chloro-2',6'-diethyl-N-(n-propoxyethyl)acetanilide, N-chloroacetyl-N-(2,6-diethylphenyl)glycine ethyl ether and 2-chloro-N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methyl)acetamide; uracil series herbicides such as 5-bromo-3-sec-butyl-6-methyluracil and 3-cyclohexyl-5,6-trimethyleneuracil; pyridinium chloride series herbicide such as 1,1'-dimethyl-4,4-bispyridinium chloride; phosphate series herbicides such as N,N-bis(phosphonomethyl)glycine, O-ethyl-O-(2-nitro-5-methylphenyl)-N-sec-butylphosphoramidothioate, S-(2-methyl-1-pyperidylcarbonylmethyl)-O,O-di-n-propyldithiophosphate and S-(2-methyl-1-pyperidylcarbonylmethyl)-O,O-diphenyldithiophosphate; toluidine series herbicide such as  $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine; 5-t-butyl-3-(2,4-dichloro-5-isopropoxyphenyl)-1,3,4-oxadiazolin-2-one; 3-isopropyl-1H-2,1,3-benzothiadiazin-(4)-3H-one-2,2-dioxide;  $\alpha$ -( $\beta$ -naphthoxy)propionanilide; 4-(2,4-dichlorobenzoyl)-1,3-dimethylpyrazol-5-yl-p-toluenesulfonate; 4'-phenylsulfonyl-1,1,1-trifluoro-

sulfono-o-toluidide; 4-chloro-5-methylamino-2-(3-trifluoromethylphenyl)pyridazin-3(2H)-one; 1-methyl-3-phenyl-5-(3-trifluoromethylphenyl)pyridin-4(1H)-one; 2-methyl-4-phenyl-sulfonyltrifluoromethylsulfoanilide; 2-(3,4-dichlorophenyl)-4-methyltetrahydro-1,2,4-oxadiazol-3,5-dione; 4-chloro-5-methylamino-2-(3-trifluoromethylphenyl)pyridazin-3(2H)-one; etc. However, the herbicides are not limited to these examples. Further, they may be also applied in combination with insecticides, nematocides, fungicides, plant growth regulators, fertilizers, etc.

Practical embodiments of the herbicidal composition according to the invention are illustratively shown in the following examples wherein parts and % are by weight. The compound number of the active ingredient corresponds to the one in Table 1.

#### Preparation Example 1

Eighty parts of Compound No. 13, 3 parts of alkyl-sulfate, 2 parts of ligninsulfonate and 15 parts of hydrous silica are well mixed while being powdered to obtain a wettable powder.

#### Preparation Example 2

Ten parts of Compound No. 22, 3 parts of alkyl-arylsulfate, 7 parts of polyoxyethylene alkylaryl ether, 60 parts of cyclohexanone and 20 parts of xylene are well mixed while being powdered to obtain an emulsifiable concentrate.

#### Preparation Example 3

0.1 Part of Compound No. 23, 1 part of hydrous

silica, 35 parts of bentonite and 63.9 parts of kaolin are well mixed while being powdered. The mixture is then kneaded with water, granulated and dried to obtain granules.

Preparation Example 4

Three parts of Compound No. 16, 0.3 part of isopropyl acid phosphate, 66.7 parts of kaolin and 30 parts of talc are well mixed while being powdered to obtain a dust.

Preparation Example 5

Twenty parts of Compound No. 25 is mixed with 60 parts of an aqueous solution containing 3 % polyoxyethylene sorbitan monolaurate and pulverized until the particle size of the active ingredient becomes less than 3 microns.

Twenty parts of an aqueous solution containing 3 % of sodium alginate as a dispersing agent are incorporated therein to obtain a suspension.

The dosage rate of the hydantoins (I) may vary on their kinds, the sorts of cultivated plants, the modes of application, etc. Generally, however, the dosage rate is from 0.05 to 50 grams, preferably from 0.1 to 20 grams, of the active ingredient per are.

The application of the hydantoins (I) as herbicides will be illustratively shown in the following Examples wherein the phytotoxicity to crop plants and the herbicidal activity on weeds were evaluated as follows: the aerial parts of the test plants were cut off and weighed (fresh weight); the percentage of the fresh weight of the treated

plant to that of the untreated plant was calculated with the latter fresh weight taken as 100; and the phytotoxicity and the herbicidal activity were evaluated by the standard given in the table below. The rating values of phytotoxicity, 0 and 1, and those of herbicidal effect; 5 and 4, are generally regarded as satisfactory to protect crop plants and to control weeds, respectively. The rating values in the paddy field test alone were calculated from the dry weight of the test plants.

Rating value	Fresh weight (percentage to untreated plot) (%)	
	Crop plant	Weeds
5	0 - 39	0
4	40 - 59	1 - 10
3	60 - 79	11 - 20
2	80 - 89	21 - 40
1	90 - 99	41 - 60
0	100	61 - 100

The following compounds were used in the Examples for comparison:


---

Compound  
No.

Structure


Remarks

(a)




British Patent  
1,503,244

(b)




U.S. patent  
3,958,976

(c)




U.S. patent  
4,179,276

(d)




U.S. patent  
4,179,276

(e)



Commercially avail-  
able herbicide known  
as "Atrazine"

(f)



Commercially avail-  
able herbicide known  
as "Ioxynil"




Compound  
No.

Structure


Remarks

(g)




Commercially available herbicide known as "Chlormethoxynil"

(h)



Commercially available herbicide known as "Alachlor"

(i)



Commercially available herbicide known as "Glyphosate"

### Test Example 1

Plastic beakers (diameter, 10 cm; height, 10 cm) were filled with upland field soil, and the seeds of barnyardgrass, wild oat, tall morningglory and velvetleaf were separately sowed in the beakers. A designed amount of the test compound formulated into an emusifiable concentrate according to Preparation Example 2 and dispersed in water was sprayed over the top by means of a small hand sprayer at a spray volume of 5 liters per are. After the spraying, the test plants were grown for 20 days in the greenhouse, and herbicidal activity was examined. The results are shown in Table 7.


0070389

Table 7

Compound No.	Dosage (weight of active ingredient, g/are)	Herbicidal activity			
		Barn-yard-grass	Wild oat	Tall morning-glory	Velvet-leaf
1	5	5	5	5	5
	2.5	5	5	4	5
2	5	5	5	5	5
	2.5	5	5	4	5
3	5	5	5	5	5
	2.5	5	5	4	5
4	5	5	5	5	5
	2.5	5	5	4	5
5	5	5	5	5	5
	2.5	5	5	4	5
6	5	5	5	5	5
	2.5	5	5	4	5
7	5	5	5	5	5
	2.5	5	5	5	5
8	5	5	5	5	5
	2.5	5	5	5	5
9	5	5	5	5	5
	2.5	5	5	4	5
12	5	5	5	5	5
	2.5	5	5	4	5
(a)	5	0	1	0	2
	2.5	0	0	0	0
(b)	5	3	1	0	5
	2.5	0	0	0	2

Test Example 2

Plastic beakers (diameter, 10 cm; height, 10 cm) were filled with upland field soil, and the seeds of barnyardgrass, wild oat, tall morningglory and velvetleaf and seeds of soybean and cotton were separately sowed in the beakers. A designed amount of the test compound formulated into an emusifiable concentrate according to Preparation Example 2 and dispersed in water was sprayed over the top by means of a small hand sprayer at a spray volume of 5 liters per are. After the spraying, the test plants were grown for 20 days in the greenhouse, and herbicidal activity and phytotoxicity were examined. The results are shown in Table 8.



0070389

- 35 -

Table 8

Compound No.	Dosage (weight of active ingredient, g/are)	Herbicidal activity				Phytotoxicity	
		Barnyard-grass	Wild oat	Tall morning-glory	Velvet-leaf	Soybean	Cotton
10	1.25	5	5	5	5	1	1
11	0.63	5	4	4	5	1	1
13	1.25	5	5	5	5	0	1
14	1.25	4	4	4	5	1	1
15	0.63	5	5	5	5	1	1
16	1.25	5	4	4	5	1	1
17	0.63	5	4	4	5	1	1
18	1.25	5	4	4	5	1	1
19	1.25	5	5	5	5	0	0
20	1.25	5	5	5	5	1	1
21	0.63	4	4	5	5	1	2
22	1.25	5	4	5	5	0	0
23	0.63	5	5	5	5	0	0
	1.25	5	5	4	5	1	1

0070389

- 36 -

(Continued)

Compound No.	Dosage (weight of active ingredient, g/are)	Herbicidal activity				Phytotoxicity	
		Barnyard-grass	Wild oat	Tall morning-glory	Velvet-leaf	Soybean	Cotton
24	1.25 0.63 1.25 0.63	5 4 5 5	5 4 4 4	4 - 5 -	5 5 5 5	- 0 - 0	- 0 - -
(c) (d)	1.25 1.25	2 4	1 3	2 3	3 4	2 3	1 3

Test Example 3

Plastic beakers (diameter, 10 cm; height, 10 cm) were filled with upland field soil, and the seeds of barnyardgrass, wild oat, wild mustard and velvetleaf were separately sowed in the beakers and grown for 2 weeks in a greenhouse. A designed amount of the test compound was sprayed to the foliage of the test plants by means of a small hand sprayer. After the spraying, the test plants were further grown for 3 weeks in the greenhouse, and herbicidal activity was examined. The results are shown in Table 9. In this treatment, the test compound was formulated into an emusifiable concentrate according to Preparation Example 2 and applied at a spray volume of 5 liters per are by dispersing it in water with the addition of a spreading agent.




Table 9


Compound No.	Dosage (weight of active ingredient, g/are)	Herbicidal activity			
		Barnyard-grass	Wild oat	Wild mustard	Velvet-leaf
1	1.25	4	4	5	5
2	5	4	4	5	5
3	1.25	5	5	5	5
4	1.25	4	3	4	5
6	5	4	4	5	5
7	1.25	5	5	5	5
8	5	4	4	5	5
13	1.25	4	5	5	5
14	1.25	3	5	5	5
15	1.25	4	4	5	5
16	1.25	5	4	5	5
18	1.25	4	4	5	5
19	1.25	4	3	5	5
20	1.25	3	3	4	5
22	1.25	4	5	5	5
(a)	5	0	0	0	5
	1.25	0	0	0	3
(b)	5	2	3	4	5
	1.25	0	1	0	3
(c)	1.25	0	0	1	4
(d)	1.25	1	1	2	5

Test Example 4

Plastic trays (35 cm x 25 cm x 15 cm) were filled with upland field soil, and the seeds of tall morningglory, velvetleaf, prickly sida, jimsonweed, black nightshade, redroot pigweed, Johnsongrass and green foxtail, and the seeds of cotton and soybean were sowed therein. A designed amount of the test compound formulated into a wettable powder according to Preparation Example 1 and dispersed in water was sprayed over the top by means of a small hand sprayer at a spray volume of 5 liters per are. After the spraying, the test plants were grown in a greenhouse for 20

0070389

days, and phytotoxicity and herbicidal activity were examined. The results are shown in Table 10.





0070389

Table 10

Com-pound No.	Dosage (weight of active ingredient, g/are)	Herbicidal activity								Phyto-toxicity	
		Tall morning-glory	Velvet-leaf	Prickly sida	Jimson-weed	Black night-shade	Redroot pigweed	Johnson-grass	Green foxtail	Cotton	Soybean
1	10	4	5	5	5	5	5	5	5	1	0
2	5	4	5	5	5	5	5	5	5	0	0
3	10	4	5	5	5	5	5	4	5	1	1
4	5	4	5	5	5	5	5	5	5	1	1
5	10	4	5	5	5	5	5	5	5	1	0
6	5	4	5	5	5	5	5	5	5	0	0
7	10	5	5	5	5	5	5	5	5	0	1
8	5	4	5	5	5	5	5	4	5	1	1
9	10	5	5	5	5	5	5	5	5	1	1
12	5	4	5	5	5	5	5	4	5	0	0
(a)	10	0	3	2	1	3	5	1	2	1	0
(b)	10	1	4	4	3	5	5	3	4	1	1
(h)	20	1	0	2	0	4	5	2	5	0	0

Test Example 5

Plastic trays (35 cm x 25 cm x 15 cm) were filled with upland field soil, and the seeds of corn, wheat, velvetleaf, cocklebur, tall morningglory, common lambsquarters, black nightshade and green foxtail were sowed and grown for 2 to 3 weeks in a greenhouse. Every two trays were placed in a frame (50 cm x 100 cm x 40 cm) and a designed amount of the test compound was sprayed thereover by means of a small hand sprayer. The test plants were further grown for 3 weeks in the greenhouse and herbicidal activity and phytotoxicity were examined. The results are shown in Table 11. In this treatment, the test compound was formulated into an emulsifiable concentrate according to Preparation Example 2 and applied by diluting it in water (25 liters) with the addition of a spreading agent. At the time of application, the plants were generally at the 1 to 4 leaf stage and in 1.5 to 20 cm height.




Table 11

Compound No.	Dosage (weight of active ingredient, g/are)	Phyto-toxicity		Herbicidal activity					
		Corn	Wheat	Velvet-leaf	Cock-lebur	Tall morning-glory	Common lambs-quarters	Black night-shade	Green fox-tail
3	0.63 0.32 0.16	1 1 0	1 1 1	5 5 5	5 5 5	5 5 4	5 5 5	5 5 5	4 2 1
4	0.63 0.32 0.16	1 1 1	1 1 1	5 5 5	5 5 5	5 5 4	5 5 5	5 5 5	3 2 4
7	0.63 0.32 0.16	1 1 1	1 1 1	5 5 5	5 5 5	5 5 5	5 5 5	5 5 5	4 3 2
8	0.63 0.32 0.16	1 1 1	1 1 1	5 5 5	5 5 5	5 5 5	5 5 5	5 5 5	4 3 2
15	0.63 0.32 0.32	1 1 0	1 1 0	5 5 5	5 5 5	5 5 5	5 5 5	5 5 5	4 4 2
19	0.63 0.32 0.32	1 0 1	0 0 0	5 5 5	5 5 5	5 5 5	5 5 5	5 4 5	4 3 5
22	0.63 0.32 0.32	0 0 1	1 1 1	5 5 5	5 5 5	5 5 5	5 5 5	5 5 5	3 4 2
25	0.63 0.32 0.32	1 0 1	1 1 1	5 5 5	5 5 5	5 5 5	5 5 5	5 5 5	4 2 2
(c)	0.63 0.32 0.63	1 1 1	1 0 1	5 3 5	4 2 4	3 2 4	3 2 5	2 1 2	1 1 1
(d)	0.63 0.32 0.63	1 1 1	1 0 1	5 3 5	4 3 5	3 3 5	3 3 1	2 2 1	1 1 1
(e)	0.63 0.32 0.32	0 1 0	1 1 1	4 5 5	3 5 3	3 5 3	1 1 1	1 1 1	0 1 0

Test Example 6

Seeds of cocklebur, common purslane, tall morningglory, jimsonweed, sicklepod and velvetleaf and the seeds of corn were sowed to the depth of 3 cm of the plotted field each having 3 m<sup>2</sup>. Eighteen days thereafter and at the time when the corn grew up to the 6-leaf stage and in 40 cm height and the weeds were at 2 to 4-leaf stages and in 5 to 15 cm height, a designed amount of the test compound was sprayed to the test plants with three replications over the top by means of a small hand sprayer. After cultivation for additional 18 days, herbicidal activity and phytotoxicity were examined. The results are shown in Table 12. In this treatment, the test compound of the invention was formulated into an emulsifiable concentrate according to Preparation Example 2 whereas that for comparison was a commercially available wettable powder. Both compounds were diluted with water containing a spreading agent and applied at a spray volume of 150 ml per plot.

---

0070389

- 44

Table 12

Compound No.	Dosage (weight of active ingredient, g/are)	Herbicidal activity							
		Phyto-toxicity	Corn	Cock-lebur	Common purslane	Tall morning-glory	Jimson-weed	Sickle-pod	Velvet-leaf
22	0.63 0.32 0.16		0 0 0	5 5 4	5 5 5	5 4 3	5 5 5	5 4 4	5 5 4
(e)	10 5 2.5		0 0 0	5 5 3	5 5 4	4 3 2	5 5 5	3 2 1	5 4 2

Test Example 7

In a plastic pot (10 cm in diameter), upland soil was filled, and tubers of purple nutsedge (Cyperus rotundus) were transplanted at the depth of 2 cm from the soil surface and cultivated in a greenhouse for 4 weeks, whereby purple nutsedge was in 7-leaved stage. A designed amount of the test compound formulated into an emulsifiable concentrate according to Preparation Example 2 and diluted with water was applied to the foliage of the test plant by means of a hand sprayer, and the test plants were further grown in the greenhouse for 8 weeks and subjected to observation by removal of the soil with water. The herbicidal activity on the aerial part (e.g. leaves) and the underground part (e.g. rhizome and tuber) was evaluated according to the same criteria as in Test Example 1. The results are shown in Table 13.




Table 13

Compound No.	Dosage of (weight of active ingredient, g/are)	Herbicidal activity	
		Aerial part	Underground part
1	20	5	5
3	10	4	4
	20	5	4
4	10	5	4
	20	5	4
7	10	5	4
	20	5	4
8	10	5	4
	20	5	4
14	10	5	4
	20	5	5
15	10	5	4
	20	5	4
22	10	5	4
	20	5	4
	10	5	4
(a)	40	2	0
(b)	40	2	1
(c)	10	1	0
(i)	40	5	4
	20	4	4

Test Example 8

Wagner's pots (1/5000 are) were filled with paddy field soil and the seeds of annual weeds (e.g. barnyard-grass, monochoria, broad-leaved weeds) were sowed to 2 to 3 cm depth, and water was poured therein until the depth of water became 4 cm. Rice seedlings of the 4-leaf stage and the tubers or buds of perenial weeds (e.g. slender spikerush, hardstem bulrush, arrowhead) were transplanted therein and grown for 5 days. At the time when the germination occured, a desinged amount of the test compound formulated in an emulsifiable concentrate according to Preparation Example 2 was applied to the pots by perfusion. Thereafter, the test plants were grown for further 4 weeks and herbicidal activity and phytotoxicity were examined. The results are shown in Table 14.







Table 14

Compound No.	Dosage (weight of active ingredient, g/are)	Herbicidal activity						Phyto-toxicity
		Barnyard-grass	Monochoria	Broad-leaved weed	Slender spikerush	Hardstem bulrush	Arrow-head	Rice plant
6	0.5 0.25	5 4	5 5	5 5	5 5	4 5	4 3	0 0
7	0.5 0.25	5 3	5 5	5 5	5 4	5 4	5 4	0 0
8	0.5 0.25	5 3	5 5	5 5	5 4	5 4	5 4	0 0
13	0.5 0.25	3 3	5 5	5 5	5 4	4 4	5 4	0 0
23	0.5 0.25	5 4	5 5	5 5	4 4	4 3	4 4	- -
(a)	0.5 0.25	1 1	3 3	3 2	1 1	1 0	1 0	0 0
(b)	0.5 0.25	2 1	3 2	3 2	2 1	1 1	1 0	0 0
(d)	0.5 0.25	1 0	2 2	2 2	2 1	1 1	0 0	0 0
(g)	0.5 0.25	2 1	2 2	2 2	1 1	1 0	1 0	0 0

What is claimed is:

1. A compound of the formula:




wherein X is a chlorine atom or a bromine atom, Y is  $-CH_2-$ ,  $-S-$  or  $-SO_2-$  and R is a  $C_1-C_4$  alkyl group, an allyl group or a propargyl group.

2. 2-(4-Chloro-2-fluoro-5-methoxyphenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine-1,3(2H,8aH)-dione.
3. 2-(4-Bromo-2-fluoro-5-methoxyphenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine-1,3(2H,8aH)-dione.
4. 2-(4-Chloro-2-fluoro-5-propoxyphenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine-1,3(2H,8aH)-dione.
5. 2-(4-Chloro-2-fluoro-5-(1-methylethoxy)-phenyl)-5,6,7,8-tetrahydroimidazo[1,5-a]pyridine-1,3(2H,8aH)-dione.
6. 2-(4-Chloro-2-fluoro-5-methoxyphenyl)-5,6,7,8-tetrahydro-1H-imidazo[5,1-c][1,4]thiazine-1,3(2H)-dione-7,7-dioxide.


7. 2-(4-Chloro-2-fluoro-5-ethoxyphenyl)-5,6,7,8-tetrahydro-1H-imidazo[5,1-c][1,4]thiazine-1,3(2H)-dione-7,7-dioxide.

8. 2-(4-Chloro-2-fluoro-5-(1-methylethoxy)-phenyl)-5,6,7,8-tetrahydro-1H-imidazo[5,1-c][1,4]thiazine-1,3(2H)-dione-7,7-dioxide.


9. A process for preparing a compound of the formula:



wherein X is a chlorine atom or a bromine atom, Y is  $-CH_2-$ ,  $-S-$  or  $-SO_2-$  and R is a  $C_1-C_4$  alkyl group, an allyl group or a propargyl group which comprises reacting a phenyl isocyanate of the formula:




wherein X and R are each as defined above with an azine of the formula:




wherein R' is an alkali metal atom or an alkyl group and Y is as defined above in an inert solvent at a temperature of

from room temperature to the boiling temperature of the solvent to give an urea the formula:




wherein X, Y, R and R' are each each as defined above, followed by treatment of the latter with a mineral acid or an alkali metal alkoxide for ring closure.

10. A process for preparing a compound of the formula:



wherein X is a chlorine atom or a bromine atom, Y is  $-CH_2-$ ,  $-S-$  or  $-SO_2-$  and R is a  $C_1-C_4$  alkyl group, an allyl group or a propargyl group which comprises reacting a hydroxyphenyl-hydantoin of the formula:




wherein X and Y are each as defined above with a halide of the formula:




wherein Z is a chlorine atom, a bromine atom or an iodine atom and R is as defined above in the presence of a base in an inert solvent at a temperature of 0 to 200°C.

11. A process for preparing a compound of the formula:



wherein X is a chlorine atom or a bromine atom and R is a C<sub>1</sub>-C<sub>4</sub> alkyl group, an allyl group or a propargyl group which comprises oxidizing a compound of the formula:



wherein X and R are each as defined above with a peracid in an inert solvent at a temperature of -30 to 30°C.


12. A herbicidal composition which comprises a herbicidally effective amount of the compound according to claim 1 as an active ingredient and an inert carrier.

13. A method for controlling weeds which comprises applying a herbicidally effective amount of the compound according to claim 1 to the area where the weeds

grow or will grow.


14. Use of the compound according to claim 1 as a herbicide.

15. A compound of the formula:



wherein X is a chlorine atom or a bromine atom and Y is  $-\text{CH}_2-$ ,  $-\text{S}-$  or  $-\text{SO}_2-$ .

16. A compound of the formula:



wherein X is a chlorine atom or a bromine atom and Y is  $-\text{CH}_2-$ ,  $-\text{S}-$  or  $-\text{SO}_2-$ .


VOSSIUS · VOSSIUS · TAUCHNER · HEUNEMANN · RAUH  
PATENTANWÄLTE  
EUROPEAN PATENT ATTORNEYS

0070389

P. O. Box 880767 - 8000 München 88

European Patent Office

8000 . Munich 2



DR. RER. NAT. VOLKER VOSSIUS  
DIPL.-CHEM. DOROTHEA VOSSIUS  
DR. RER. NAT. PAUL TAUCHNER  
DR. RER. NAT. DIETER HEUNEMANN  
DR. RER. NAT. PETER RAUH


SIEBERTSTRASSE November 3, 1982  
P. O. BOX 880767 DV/ge  
8000 MÜNCHEN 88  
PHONE: (089) 47 4075  
CABLE: BENZOLPATENT MÜNCHEN  
TELEX: 5-29453 VOPAT D

Re.: EP 82105237.0  
Sumitomo Chemical Company, Limited

OUR REF.:  
UNSER ZEICHEN: R 922 EP


In reply to the Communication of September 20, 1982:

There is an obvious error in formula (f) on page 31. It is hereby requested to change formula (f)



"Ioxynil"

to the correct formula



"Ioxynil"

*D. Vossius*  
Dorothea Vossius  
Patentanwältin

The request for correction is allowed under  
R. 56 EPC / with the exception of the deleted  
part.  
15 NOV 1982  
RECEIVING SECTION  
A. G. POELS

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**